


Nonlocal Quantum Kinetic Theory



MAX-PLANCK-GESSELLSCHAFT

Real-time Green functions

Two independent correlation functions $1 = r, t, s, \dots$
 $G^>(1, 2) = < a(1)a^+(2) >$ and $G^<(1, 2) = < a^+(2)a(1) >$

Martin-Schwinger hierarchy for causal GF
 $iG = \Theta(t_1 - t_2)G^> \pm \Theta(t_2 - t_1)G^<$

$$G_0^{-1}G(1, 1') = \delta_{1,1'} + \int d3V(1, 3)G_2(1, 3, 1', 3') \equiv \delta_{1,1'} + \int_c d3\Sigma(1, 3)G(3, 1')$$

$$\text{with } G_0^{-1}(11') = \left(i \frac{\partial}{\partial t_1} - \frac{(\frac{\partial}{\partial t_1})^2}{2m} - \Sigma^{HF}(11') \right) \delta(1 - 1')$$

Weakening of initial correlations $\lim_{t \rightarrow t_0} \Sigma G = 0$ leads to Keldysh contour

$$\int_c d\bar{1}\Sigma(1, \bar{1})G(\bar{1}, 1') = \int_{-\infty}^{+\infty} d\bar{1} \{ \Sigma(1, \bar{1})G(\bar{1}, 1') - \Sigma^<(1, \bar{1})G^>(\bar{1}, 1') \}$$

Consequence: Kadanoff and Baym equation (KB)

$$-i(G_0^{-1}G^< - G^<G_0^{-1})(t_1, t_2) = \int_{t_0}^{t_1} \Sigma^>G^< + \int_{t_0}^{t_2} G^<\Sigma^> - \int_{t_0}^{t_1} G^>\Sigma^< - \int_{t_0}^{t_2} \Sigma^<G^> + \int_{t_1}^{t_2} [\Sigma^<, G^>]$$

Gradient expansion of KB -equation

First order gradient expansion $t_1 - t_2 \rightarrow \omega, r_1 - r_2 \rightarrow k$

$$\left(\frac{\partial}{\partial t} + \left(\frac{p}{m} + \nabla_p \Sigma^{HF} \right) \nabla_R - \nabla_R \Sigma^{HF} \nabla_p \right) \rho = \int \frac{d\omega}{2\pi} (G_\omega^>\Sigma_\omega^< - G_\omega^<\Sigma_\omega^>) + \int \frac{d\omega d\omega'}{(2\pi)^2} \frac{P}{\omega' - \omega} (\{\Sigma_\omega^>, G_\omega^<\} - \{\Sigma_\omega^<, G_\omega^>\}) - \frac{\partial}{\partial t} \int \frac{d\omega' d\omega}{(2\pi)^2} \frac{P'}{\omega' - \omega} (G_\omega^>\Sigma_\omega^< - G_\omega^<\Sigma_\omega^>).$$

leads to quasiparticle distribution and iteration for Wigner function

$$\rho = \int \frac{d\omega}{2\pi} G_\omega^< \approx f - \int \frac{d\omega'}{2\pi} \frac{P'}{\omega' - \varepsilon} ((1 - f)\Sigma_\omega^< - f\Sigma_\omega^>)$$

kinetic equation for quasiparticles

$$\frac{\partial}{\partial t} f + \nabla_k \varepsilon \nabla_R f - \nabla_R \varepsilon \nabla_k f = z((1 - f)\Sigma^< - f\Sigma^>)$$

with wave function renormalization $z = (1 - \partial_\omega \Sigma)^{-1}$

$$\rho = z f - \int \frac{d\omega}{2\pi} \frac{P'}{\omega - \varepsilon} \Sigma_\omega^< \quad \text{corresponds to extended quasiparticle picture}$$

$$A = G^> - G^< = \frac{\Gamma_\omega}{[\omega - \frac{p^2}{2m} - \Sigma_\omega]^2 + \frac{1}{4}\Gamma_\omega^2} \approx 2\pi z\delta(\omega - \varepsilon) - \Gamma_\omega \frac{\partial}{\partial \omega} \frac{\rho}{\omega - \varepsilon}$$


Equilibrium:
 Craig '66, Bezerides DeBois '68, Zimmermann, Stoltz '79, Kremp '84, Röpke, Schmidt '87, Köhler, Malfliet '93

Nonequilibrium:
 Lipavský, Špička 1995, Morawetz '00

Message from Equilibrium

$$\rho(k) = \int \frac{d\omega}{2\pi} \frac{\Gamma(\omega, k)}{(\omega - \frac{p^2}{2m} - \Sigma(\omega, k))^2 + \frac{1}{4}\Gamma(\omega, k)^2} f_{FD}(\omega)$$


example from nuclear matter



Spectral function for selfconsistent separable T-matrix at $T = 10\text{MeV}$

Full spectral function $k = 0.7 \text{ fm}^{-1}$, free particle, Hartree Fock.

Extended Quasiparticle Picture $\rho = f + \int \frac{d\omega}{2\pi} (\Sigma^<(1 - f) - \Sigma^>f) \frac{P'}{\omega - \varepsilon}$



Non-local kinetic equation

Non-Markovian (memory) kinetic equation for reduced density ρ

$$\frac{\partial}{\partial t} \rho + \nabla_k \epsilon_{HF} \nabla_R \rho - \nabla_R \epsilon_{HF} \nabla_k \rho = \int_0^{t-t_0} d\tau \left(\{G^<(t-\frac{\tau}{2}, \tau), \Sigma^>(t-\frac{\tau}{2}, -\tau)\} - \{G^>(t-\frac{\tau}{2}, \tau), \Sigma^<(t-\frac{\tau}{2}, -\tau)\} \right)$$

is transformed by extended quasiparticle picture into precursor of kinetic equation for quasiparticle distribution f

$$\frac{\partial}{\partial t} f + \nabla_k \epsilon \nabla_R f - \nabla_R \epsilon \nabla_k f = z((1 - f)\Sigma^< - f\Sigma^>)$$

ladder summation: Boltzmann-Uehling-Uhlenbeck, random phase approximation: Lenard- Balescu equation, etc

But: non-local scattering events by $\Sigma[G] \rightarrow \Sigma[f]$

Memory or off-shell parts

- compensate the off-shell parts in Kadanoff Baym equation without other neglects
- in quasiparticle energy $\varepsilon_1 = \frac{k^2}{2m_a} + \text{Re}\Sigma_{1,\varepsilon_1}^R$, off-shell part leads to the correct binding energy
- Wave function renormalization z
- A genuine time nonlocality Δ_t
- direct link between Wigner function (reduced density matrix) and quasiparticle distribution

virial corrections from intrinsic gradients in the scattering integrals.

Consistent kinetic equation

$$\frac{\partial f_1}{\partial t} + \frac{\partial \varepsilon_1 \partial f_1}{\partial k} - \frac{\partial \varepsilon_1 \partial f_1}{\partial r} = \int \frac{dp dq}{(2\pi)^5} \mathcal{P} \delta(\varepsilon_1 + \varepsilon_2^- - \varepsilon_3^- - \varepsilon_4^- - 2\Delta_E) \times \left[(1-f_1)(1-f_2^-)f_3^-f_4^- - f_1f_2^-(1-f_3^-)(1-f_4^-) \right]$$

where

$$\begin{aligned} f_1 &\equiv f(k, r, t) \\ f_2^- &\equiv f(p, r - \Delta_2, t) \\ f_3^- &\equiv f(k - q - \Delta_3, r - \Delta_3, t - \Delta_t) \\ f_4^- &\equiv f(p + q - \Delta_4, r - \Delta_4, t - \Delta_t) \end{aligned}$$

with T-matrix $\boxed{T} = \boxed{+} + \boxed{T} = |\mathbf{T}| e^{i\phi}$


$$\mathcal{P} = z_1 z_2 z_3 z_4 |\mathbf{T}|^2 (\varepsilon_1 + \varepsilon_2^- - \Delta_E, k - \frac{\Delta_K}{2}, p - \frac{\Delta_K}{2}, q, r - \frac{1}{4}(\Delta_2 + \Delta_3 + \Delta_4), t - \frac{\Delta_t}{2})$$

$$\Delta_t = \frac{\partial \phi}{\partial t} \Big|_{\varepsilon_1 + \varepsilon_2} \quad \Delta_2 = \left(\frac{\partial \phi}{\partial p} - \frac{\partial \phi}{\partial q} - \frac{\partial \phi}{\partial k} \right) \Big|_{\varepsilon_1 + \varepsilon_2}$$

$$\Delta_E = -\frac{1}{2} \frac{\partial \phi}{\partial t} \Big|_{\varepsilon_1 + \varepsilon_2} \quad \Delta_3 = -\frac{\partial \phi}{\partial k} \Big|_{\varepsilon_1 + \varepsilon_2}$$

$$\Delta_K = \frac{1}{2} \frac{\partial \phi}{\partial k} \Big|_{\varepsilon_1 + \varepsilon_2} \quad \Delta_4 = -\left(\frac{\partial \phi}{\partial k} + \frac{\partial \phi}{\partial q} \right) \Big|_{\varepsilon_1 + \varepsilon_2}$$

P. Lipavský, K. M., and V. Špička: *Kinetic equation for strongly interacting dense Fermi systems*, Annales de physique, 26, 1 (2001) ISBN 2-86883-541-4



Energy conversion: latent heat

There appear an internal energy- $\mathcal{E}_{\text{gain}} = \int d\mathcal{P} \Delta_E$ and momentum-gain $\mathcal{F}_{\text{gain}} = \int d\mathcal{P} \Delta_K$

From nonlocal kinetic equation

$$\int \frac{dk}{(2\pi)^3} \varepsilon_k \frac{\partial f_k}{\partial t} = -\frac{d}{dt} \int d\mathcal{P} \frac{\varepsilon_k + \varepsilon_p}{2} \Delta_t + \int d\mathcal{P} \Delta_E$$

energy gain combines together with drift term into time derivative

$$\sum_a \int \frac{dk}{(2\pi)^3} \varepsilon \frac{\partial f}{\partial t} - \mathcal{E}_{\text{gain}} = \frac{\partial \mathcal{E}_{\text{qp}}}{\partial t}.$$

of the quasiparticle energy functional

$$\mathcal{E}_{\text{qp}} = \sum_a \int \frac{dk}{(2\pi)^3} f_a(k) \frac{k^2}{2m} + \frac{1}{2} \sum_{ab} \int \frac{dk dp}{(2\pi)^6} f_a(k) f_b(p) \mathcal{T}_{\text{ex}}(\varepsilon_1 + \varepsilon_2, k, p, 0)$$

• By energy gain $\Delta_E = \frac{\partial \mathcal{E}}{\partial t}$ transformation of kinetic energy into correlation energy. Similar to breathing hard-sphere by Pauli-blocking

Entropy balance

$$\frac{\partial(\mathcal{S}^{\text{qp}} + \mathcal{S}^{\text{mol}})}{\partial t} + \frac{\partial(j^{\text{qp}} j^{\text{mol}})}{\partial r} = \mathcal{T}_{\text{gain}}^S$$

quasiparticle part $\mathcal{S}^{\text{qp}} = -k_B \int \frac{dk}{(2\pi)^3} [f_1 \ln f_1 + (1 - f_1) \ln (1 - f_1)]$
 molecular part $\mathcal{S}^{\text{mol}} = -\frac{k_B}{2} \int d\mathcal{P} f_1 f_2 (1 - f_3 - f_4) \ln \frac{f_3 f_4}{(1 - f_3)(1 - f_4)}$
 entropy gain

$$\mathcal{T}_{\text{gain}}^S = -\frac{k_B}{2} \sum_{ab} \int d\mathcal{P} f_1 f_2 (1 - f_3) (1 - f_4) \ln \frac{f_3 f_4 (1 - f_1) (1 - f_2)}{(1 - f_3) (1 - f_4) f_1 f_2} \geq 0$$

• H-theorem holds also for nonlocal kinetic theory

Relation to Landau theory

Landau theory works only if collisions \tilde{I} treated instant and local
 From kinetic equation $\frac{\partial f_k}{\partial t} = \tilde{I}_k$ follows:

number of particles

$$\frac{dn}{dt} = \int \frac{dk}{(2\pi)^3} \frac{\partial f_k}{\partial t} = \int \frac{dk}{(2\pi)^3} \tilde{I}_k \equiv 0$$

$$\frac{d\mathcal{E}}{dt} = \int \frac{dk}{(2\pi)^3} \frac{\partial \mathcal{E}}{\partial k} \frac{\partial \tilde{I}_k}{\partial t} = \int \frac{dk}{(2\pi)^3} \tilde{e}_k \tilde{I}_k \equiv 0$$

Nonlocal kinetic theory

$$\frac{dn}{dt} = \frac{d}{dt} \int \frac{dk}{(2\pi)^3} \tilde{I}_k + \frac{d}{dt} \int d\mathcal{P} \Delta_t \quad \frac{d\mathcal{E}}{dt} = \int \frac{dk}{(2\pi)^3} \varepsilon_k \frac{\partial \tilde{I}_k}{\partial t} + \frac{d}{dt} \int d\mathcal{P} \frac{\varepsilon_k + \varepsilon_p}{2} \Delta_t - \int d\mathcal{P} \Delta_E$$

Instant approximation, last term can be rewritten $\Delta_E = -\frac{1}{2} \frac{\partial \phi}{\partial t}$

$$-\int d\mathcal{P} \Delta_E = \frac{1}{2} \int d\mathcal{P} \frac{\partial \phi}{\partial t} = \int \frac{dk}{(2\pi)^3} \int d\mathcal{P} \frac{\partial \phi}{\partial k} \frac{\partial \tilde{I}_k}{\partial t} \equiv \int \frac{dk}{(2\pi)^3} \varepsilon_k \frac{\partial \tilde{I}_k}{\partial t}$$

with rearrangement energy
 follows variational expression of Landau theory $\tilde{e}_k = \varepsilon_k + \frac{\Delta}{k}$
 Landau theory mimics for energy gain, but no correlated density !

Summary

1. Non-local collisions in quantum kinetic equation
 - Unifies Landau - quasiparticle and Enskog - like equations
 - Thermodynamically consistent equation of state (energy, density, pressure) including quantum 2. virial coefficient
 - Quantum correlation recast into quasi-classical picture
 - Particle-hole vs space-time symmetry completed
 - Consistent theory unifying Landau theory and dense gases
 - Explicit calculation of Wigner function not necessary, correlated observables directly from nonlocal kinetic equation
 - No additional computational expenses
2. Memory effects result in
 - Off-shell tails of Wigner function (exactly compensated)
 - Renormalization of scattering rates
 - Collision delay is all what is left from memory
3. Successfully applied to: Heavy ion collisions, deep neutral impurities, Bernoulli potential at superconducting surfaces, ...

Literature:

<http://www.pks.mpg.de/~morawetz>

I. Essentials

-Kinetic equation for strongly interacting dense Fermi systems, Annals of Physics, Vol. 26 No. 01 (February